


RESIDUAL MAGNETIC ANOMALY AND FREE-AIR GRAVITY ANOMALY
PROFILES, 1973, ON CONTINENTAL SHELF AND SLOPE BETWEEN BERING
STRAIT AND BARROW, ALASKA, AND MACKENZIE BAY, CANADA

BY
WILLIAM F. HANNA, BYRON D. RUPPEL, A. CRAIG McHENDRIE, AND
ROBERT F. SIKORA


Plate 3. Profiles for Lines 303 to 314. Horizontal axes are marked in units of hours and kilometers (approximate). Times are referenced to Alaskan Daylight Time (ship time), nine hours earlier than Greenwich Mean Time. Geophysical data obtained using a proton-precession magnetometer and a ship-borne gravimeter. Residual magnetic anomalies obtained by subtracting the International Geomagnetic Reference Field from the observed magnetic field.

U.S. Geological Survey OpenFile Map
Map released without editorial and
technical review for conformity to
U.S. Geological Survey standards


LINE 303-307



LINE 309



LINE 311-313



FREE AIR ANOMALY-X RESIDUAL MAGS-+

FREE AIR ANOMALY-X RESIDUAL MAGS-+

FREE AIR ANOMALY-X RESIDUAL MAGS-+